

ARTÍCULO ORIGINAL

## Evaluation of the relationship between self-reported physical activity and metabolic syndrome and its components in apparently healthy women

Milton Fabián Suárez-Ortegón<sup>1</sup>, Alejandra Arbeláez<sup>1</sup>, Mildrey Mosquera<sup>1</sup>,  
Robinson Ramírez-Vélez<sup>2</sup>, Cecilia Aguilar-De Plata<sup>1</sup>

<sup>1</sup> Grupo de Nutrición, Departamento de Ciencias Fisiológicas, Universidad del Valle, Cali, Colombia

<sup>2</sup> Postgrados de Rehabilitación, Escuela de Medicina y Ciencias de la Salud, Universidad del Rosario, Bogotá, D. C., Colombia

**Introduction:** The metabolic syndrome, a set of metabolic anomalies that include insulin resistance, central obesity, dyslipidemia, hypertension and inflammation, is an important tool to explore factors associated to cardiometabolic disease.

**Objective:** The aim of this study was to evaluate the relationship of the levels of self-reported physical activity and the International Physical Activity Questionnaire items and the metabolic syndrome and the variables related to cardiovascular risk in 89 women.

**Materials and methods:** The short version of International Physical Activity Questionnaire was applied to classify participating subjects into three categories: insufficient, sufficient and very active physical activity. The metabolic syndrome was assessed according to the International Diabetes Federation criteria. Biochemical and anthropometrical parameters were measured.

**Results:** Twenty-two participants (23%) presented metabolic syndrome and 66 women (74.2%) were classified in the insufficient physical activity category. No association was found between insufficient physical activity and metabolic syndrome. Inverse correlations were found among the days and minutes per week of physical activity of moderate-intensity, waist circumference ( $\rho=-0.327$ , and  $\rho=-0.313$ ,  $p<0.005$ , respectively), and body mass index ( $\rho=-0.262$ , and  $\rho=-0.218$ ,  $p<0.05$ , respectively).

**Conclusion:** A high prevalence of insufficient physical activity was found in the study participants, but this was not associated with metabolic syndrome. Moderate but not vigorous physical activity items from the International Physical Activity Questionnaire correlated inversely with anthropometrical markers related to cardiovascular risk.

**Key words:** Motor activity, women, cardiovascular diseases.

doi: <http://dx.doi.org/10.7705/biomedica.v34i1.1442>

### Evaluación de la relación de actividad física autorreportada con el síndrome metabólico y sus componentes en mujeres aparentemente sanas

**Introducción.** El síndrome metabólico, conjunto de anomalías metabólicas que incluyen resistencia a la insulina, obesidad central, dislipidemia, hipertensión e inflamación, es una herramienta importante para explorar los factores asociados a enfermedades cardiovasculares.

**Objetivo.** El objetivo de este estudio fue evaluar la relación de los niveles autorreportados de actividad física y los elementos del Cuestionario Internacional de Actividad Física (*International Physical Activity Questionnaire*), con el síndrome metabólico y las variables relacionadas con el riesgo cardiovascular en 89 mujeres.

**Materiales y métodos.** La versión corta del Cuestionario Internacional de Actividad Física se aplicó para clasificar a los sujetos en tres categorías: actividad física insuficiente, suficiente y muy activa. El síndrome metabólico se evaluó según los criterios de la Federación Internacional de Diabetes y se midieron los parámetros bioquímicos y antropométricos.

**Resultados.** Veintidós participantes (23 %) presentaron síndrome metabólico y 66 mujeres (74,2 %) fueron clasificadas en la categoría de actividad física insuficiente. No se encontró asociación entre la actividad física insuficiente y el síndrome metabólico. Se encontraron correlaciones inversas entre los días y minutos de actividad física de moderada intensidad por semana con la circunferencia de cintura

#### Author contributions:

Milton Fabián Suárez-Ortegón participated in designing the project, in field work, statistical analysis and in drafting the document. Alejandra Arbeláez, Mildrey Mosquera, Cecilia Aguilar-De Plata participated in designing the project, in field work and in correcting the document.

Robinson Ramírez-Vélez participated in drafting the document.

( $\rho=-0,327$ , y  $\rho=-0,313$ ,  $p<0,005$ , respectivamente) y el índice de masa corporal ( $\rho=-0,262$ , y  $\rho=-0,218$ ,  $p<0,05$ , respectivamente).

**Conclusión.** Una alta prevalencia de actividad física insuficiente se encontró en las participantes en el estudio, pero esto no se asoció con el síndrome metabólico. Los elementos de la actividad física moderada del Cuestionario Internacional de Actividad Física, pero no así los de actividad vigorosa, se correlacionan inversamente con marcadores antropométricos relacionados con riesgo cardiovascular.

**Palabras clave:** actividad motora, mujeres, enfermedades cardiovasculares.

doi: <http://dx.doi.org/10.7705/biomedica.v34i1.1442>

Cardiometabolic diseases constitute the main cause of world mortality and represent almost 60 % of all deaths and 43 % of all disease in the world (1). During the 1980s, Reaven, *et al.* (2), observed that dyslipidemia, hypertension and hyperglycemia were events frequently associated in the same individual, a condition that increased cardiovascular risk and became known as the metabolic syndrome. Metabolic syndrome is a set of metabolic anomalies that include glucose intolerance, insulin resistance, central obesity, atherogenic dyslipidemia, hypertension, inflammation, and prothrombotic state. This metabolic alteration, initially described by Reaven (2), has been found in subjects who report conducts and preferences related to unhealthy lifestyles, which include inadequate diet, sedentary habits, alcohol consumption and smoking.

According to the National Health and Nutrition Surveys, the current obesity epidemic and the high levels of sedentary habits have duplicated the prevalence of metabolic syndrome in American overweight adults during a period of only 10 years (3,4). The prevalence of metabolic syndrome in the USA (NHANES III Study) (5), found using the diagnosis criteria from the Third Panel of Experts from the National Cholesterol Education Program (NCEP-ATP III) (6), was close to 24% in 20-year-old and older individuals. Combined efforts by the International Diabetes Federation (IDF) (7), the US National Heart, Lung, and Blood Institute (NHLBI) and the American Heart Association (AHA) produced a new definition of metabolic syndrome to be used in clinical practice.

This health problem is becoming a critical issue in developing countries highly influenced by modernization and urbanization (8). Lifestyle

changes including reduction in physical activity and substitution of traditional diets for diets with high fat content and caloric density are some of the explanations of this phenomenon (9). Additionally, age, gender, and genetics are factors with significant impact on the predisposition to metabolic syndrome.

Several studies have evaluated the association among cardiovascular risk factors, metabolic syndrome, and the different physical activity characteristics defined by the standards of the International Physical Activity Questionnaire (IPAQ) with uncertain results (10,11). The objective of this study was to evaluate the relationship between the levels of self-reported physical activity and IPAQ items and metabolic syndrome and other variables related to cardiovascular risk in 89 women from the urban area of Cali, Colombia.

### Materials and methods

The study included 89 women workers from the public and private sectors aged between 25 and 64 years from whom blood samples were taken shortly after the interview and after having obtained informed signed consent. Exclusion criteria included women with medical or clinical diagnosis of systemic disease (including malignant processes), non-transmissible chronic disease (such as diabetes, hypertension, obesity, dyslipidemia, atherosclerosis), a history of drug or alcohol abuse, intake of multivitamins, use of statins at the moment of inclusion in the study, pregnancy, current or previous breastfeeding (<1 year), and inflammation disorders or systemic infection. This study was approved by the institutional Human Ethics Committee of the Universidad del Valle.

A trained survey taker applied the self-reported short version of the IPAQ recommended by the World Health Organization (WHO) as a valid measurement to estimate physical activity. This version has seven questions regarding the frequency, duration, and intensity of participation

Corresponding author:

Milton Fabián Suárez-Ortegón, Universidad del Valle (San Fernando), Calle 4B N° 36-00, edificio 116, quinto piso, oficina 5503, Cali, Colombia  
Tel.: (572) 518 5603  
[fabian.suarezuv@gmail.com](mailto:fabian.suarezuv@gmail.com)

Recibido: 29/10/12; aceptado: 22/08/13

in physical activity (such as walking or running during the week) immediately before participating in the study and in different daily life contexts. Participants were classified based on their weekly energy expenditure expressed as metabolic equivalents (MET):  $\leq 600$  MET/wk, insufficiently active; 601-1,500 MET/wk, sufficiently active, and  $\geq 1.500$  MET/wk, very active (12).

Waist circumference was measured between the lower rib and the iliac crest with a measuring tape; participants were asked to remain in standing position and undressed. Body mass index (BMI) was calculated as weight/height<sup>2</sup>. Body composition was determined through bio-impedance analysis using the Bodystat® device (Quadscan 4000, UK) to indirectly calculate body fat percentage from total fat mass (kg) and body weight.

Blood pressure was measured with a digital sphygmomanometer (OMRON®) on the right arm on two different occasions, five minutes apart, with the participants seated comfortably and after ten minutes of rest.

Via antecubital venipuncture, 10 ml of blood were extracted onto vacutainer tubes without additive. The blood samples were taken to the laboratory in less than one hour in coolers kept between 4 and 8 °C, and they were centrifuged at 3,000 rpm to obtain serum until processing. The biochemical markers were determined as follows: lipid levels and glycemia, via semi-automated A-15 colorimetric assay (Biosystems, Spain).

The study followed the International Diabetes Federation criteria and definition of metabolic syndrome. It was necessary to evaluate the presence of abdominal obesity (waist circumference  $\geq 80$  cm), triglycerides ( $\geq 150$  mg/dl), low levels of HDL-c ( $\leq 50$  mg/dl), systolic blood pressure ( $\geq 130$  mm Hg), diastolic blood pressure ( $\geq 85$  mm Hg), and fasting glycemia ( $\geq 100$  mg/dl) (13).

An exploratory analysis was conducted to determine the frequency and distribution of each of the variables studied. We used the Spearman correlation to estimate the relationship between the IPAQ items and variables related to cardiovascular risk. The association of the lowest physical activity level with metabolic syndrome and its components was estimated by logistic regression. A value of  $p < 0.05$  was considered as significant and all the analyses were performed with the SPSS statistical package (Statistical Program, version 13, Chicago, IL).

## Results

In table 1, the means and standard deviations of the study variables are shown. Metabolic syndrome prevalence and insufficient physical activity were 10.1 and 74.6%, respectively.

### Metabolic syndrome and insufficient physical activity

Table 2 describes the odds ratio (OR) for metabolic syndrome and its components after adjusting the insufficient physical activity level (vs. the other levels together) by age and menopause. The women reporting the lowest level of physical activity presented a 2.53-fold increased risk of having MS, but the confidence interval was not significant. The OR between insufficient physical activity level and metabolic syndrome components was also not significant.

**Table 1.** Clinical, anthropometric and biochemical variables related to cardiovascular risk, metabolic syndrome and physical activity level in urban area women

N	89
<b>Age (years)<sup>1</sup></b>	47.1 $\pm$ 7.7
<b>Blood pressure (mm Hg)<sup>1</sup></b>	
Systolic	115 $\pm$ 19
Diastolic	73 $\pm$ 14
<b>Anthropometrics<sup>1</sup></b>	
Waist circumference (cm)	77.3 $\pm$ 8.2
Body mass index (kg/m <sup>2</sup> )	26.8 $\pm$ 5.2
Body fat (%)	39.6 $\pm$ 5.8
<b>Biochemical<sup>1</sup></b>	
Glucose (mg/dl)	86.2 $\pm$ 8.8
Triglycerides (mg/dl)	125.2 $\pm$ 57.4
Cholesterol (mg/dl)	194.2 $\pm$ 36.9
c-HDL (mg/dl)	51.9 $\pm$ 10.5
<b>Metabolic syndrome, n (%)</b>	
<b>Physical activity level (MET/wk), n (%)</b>	
Insufficiently active (<600)	9 (10.1)
Sufficiently active (601 to 1,500)	66 (74.2)
Very active (>1,500)	16 (18.0)
	7 (7.9)

<sup>1</sup> Average  $\pm$  standard deviation

**Table 2.** Odd ratios<sup>1</sup> of metabolic syndrome and its components by insufficient physical activity (vs. the other levels)

	Odd ratios	95% CI
Metabolic syndrome	2.53	0.28-22.2
Systolic blood pressure $\geq 130$ mm Hg and/or diastolic $\geq 85$ mm Hg	1.15	0.32-4.10
Triglycerides $> 150$ mg/dl	0.71	0.22-2.25
c-HDL $< 50$ mg/dl	2.25	0.78-2.47
Glucose $> 100$ mg/dl	1.18	0.10-13.53
Waist circumference $> 80$ cm	1.15	0.38-3.44

<sup>1</sup> Adjusted for age and menopause  
CI: confidence interval

### **International Physical Activity Questionnaire items and variables related to cardiovascular risk**

Significant inverse correlations were found between days and time (in days and minutes per week) of moderate physical activity and the anthropometric markers waist circumference and BMI (table 3). There were no statistically significant correlations between total physical activity (MET/week) and clinical, anthropometric and biochemical variables (data not shown).

### **Discussion**

The objective of this study was to evaluate the relationship between different levels of physical activity and metabolic syndrome risk factors in 89 women from the urban area of Cali, Colombia. We used the IPAQ short version, a measuring instrument endorsed by the WHO, to estimate the level of physical activity in a population between 15 and 69 years of age. The questionnaire psychometric properties, as well as its validity and reproducibility, are applicable for prevalence studies based on national populations (14,15). Currently, the IPAQ is broadly known as the most objective way of estimating physical activity level (16-18). The assessment of physical aptitude has been considered relevant for health status because physical activity is a protection factor in relation to the development of chronic non-transmissible diseases, mainly coronary disease and metabolic syndrome (19). However, there are different factors that can modify this indicator. It is suggested that age is a factor associated with a decrease in physical activity, and this is attenuated positively in trained individuals who participate in the habitual practice of physical

exercise (20,21). An epidemiological study carried out in the 1980s by Heath, *et al.* (22), supported this hypothesis by showing that physical aptitude estimated by the level of physical activity performed diminished by 9% per decade in sedentary subjects while in trained subjects, the reduction was only in the order of 5% per decade.

With regard to the general prevalence of subjects with insufficient physical activity in this study, the value was 72.6%, which is 2.65-fold higher than that recently described by Rodrigues, *et al.* (23), and Martins, *et al.* (24), as well as higher than that described in other studies in Colombia (25,26). However, the reasons for this high prevalence were not evaluated. A factor that may have contributed to the increased proportion of sedentary activities is related to the aspects of the time available for this purpose. In fact, when analyzing the motives for physical inactivity in other works, the main reason was insufficient time (27,28). Similar reasons were found by Marcondelli, *et al.* (29), in a comparable population to the one participating in this study, with 66.7% of those surveyed reporting lack of time. However, this motive does not oppose that of increasing daily physical activity through aerobic activities, even moderate ones such as walking from three to five times per week for 30 minutes per session, which can improve cardiovascular health (30). Consequently, lack of time is a possible factor limiting physical activity.

Although many studies have found a relationship between self-reported physical activity level and metabolic syndrome (31-33), others, such as the present study, have not observed any association

**Table 3.** Spearman correlations of International Physical Activity Questionnaire survey items with clinical, anthropometric and biochemical variables related to cardiovascular risk

	Walking (min/week)	Intense physical activity (min/week)	Moderate physical activity (min/week)	Walking (days/week)	Intense physical activity (days/week)	Moderate physical activity (days/week)	Moderate physical activity (min/week)
<b>Blood pressure (mm Hg)</b>							
Systolic	0.147	0.033	-0.135	0.116	0.041	-0.092	-0.096
Diastolic	0.069	-0.025	-0.163	0.077	-0.012	-0.132	-0.042
<b>Anthropometrics</b>							
Waist circumference (cm)	0.107	-0.152	<b>-0.313**</b>	-0.025	-0.161	<b>-0.327**</b>	0.021
Body mass index (kg/m <sup>2</sup> )	0.154	-0.095	<b>-0.218*</b>	0.059	0.088	<b>-0.262*</b>	0.057
Body fat (%)	0.037	-0.101	-0.153	0.032	0.099	-0.184	0.040
<b>Biochemicals</b>							
Glucose (mg/dl)	<b>0.231*</b>	-0.188	-0.029	0.141	-0.191	-0.029	-0.061
Triglycerides (mg/dl)	0.041	0.004	-0.090	0.014	-0.014	-0.066	-0.025
Cholesterol (mg/dl)	0.002	-0.001	-0.037	-0.005	-0.026	-0.036	0.069
c-HDL (mg/dl)	-0.030	0.131	-0.007	0.015	0.108	-0.018	-0.056

\*  $p < 0.05$ , \*\*  $p < 0.005$

(34-36) (table 4). Dalacorte, *et al.*, attributed the lack of relationship to the long time required to establish a disorder such as metabolic syndrome and the possibility that numerous individuals with metabolic syndrome may have been sufficiently active and very active recently (33). Even a study with a bigger sample size than that of the study by Dalacorte, *et al.*, did not find a relationship between self-reported physical activity level and metabolic syndrome (34). On the other hand, the women participating in the present study were apparently healthy, without clinical conditions such as hypertension or diabetes, which could represent a bias with respect to regular physical activity as a part of treatment in this situation.

The IPAQ items related to moderate physical activity (time and days per week) and walking (time per week) were the only survey parameters significantly correlated with some variables related to cardiovascular risk. In the case of time and days of moderate physical activity, the correlations were exclusively with the anthropometric markers waist circumference and BMI, but in an inverse way. These inverse correlations are coherent with the premise of reduction in the recognized predictors of cardiovascular disease by regular practice of physical activity. However, time and days per week of vigorous intensity physical activity did not correlate

with clinical, anthropometrical, and biochemical variables. In a prospective study, Wannamethee and Shaper described the association between moderate intensity physical activity and significant reduction in the risk of stroke and heart attacks in men both with and without pre-existing ischemic heart disease; however, vigorous activities were not associated (37). In contrast, Hu, *et al.*, reported a reduction in the risk of type 2 diabetes through both vigorous and moderate intensity physical activity (38). Moreover, the correlation between walking minutes per week and basal glucose levels was unexpectedly positive. However, in this correlation, age could be a confounding factor because this variable correlated positively with basal glucose levels ( $\rho=0.334$ ,  $p=0.001$ ) and walking minutes per week ( $\rho=0.231$ ,  $p=0.029$ ). In this case, it is possible that older women tend to practice more walking and have higher glucose levels, or that this unexpected finding between fasting glucose and walking minutes per week could be a simple circumstantial and statistical relationship.

Some limitations must be considered in this study, one being the fact that ethnicity was not an adjustment variable. Another aspect to highlight is the measurement and quality of the responses to the IPAQ, which could be deficient (overestimating or underestimating the level of physical activity).

**Table 4.** Findings regarding the association between self-reported physical activity and metabolic syndrome in some studies

Country, authors (Reference)	n	Questionnaire	Metabolic syndrome criteria	Association
China Yu, <i>et al.</i> (31)	1,831 women/ 1,458 men	IPAQ, short form	ATP III	Yes. Higher levels of total physical activity had a lower risk of having metabolic syndrome.
Iran Esteghamati, <i>et al.</i> (32)	1,629 women/ 1,667 men	GPAQ (Global Physical Activity Questionnaire)	ATP III	Yes. Highest metabolic syndrome prevalence in individuals with lowest physical activity level
Líbano Sibai, <i>et al.</i> (33)	284 women/ 215 men	IPAQ, short form	IDF	Yes. Lack of physical exercise was associated significantly with higher odds ratio for metabolic syndrome.
Brazil Dalacorte, <i>et al.</i> (34)	246 women/ 116 men	IPAQ, short form	IDF	No. Odds ratio not significant for metabolic syndrome presence by physical activity levels
Irán Sann, <i>et al.</i> (35)	984 women	IPAQ, long form	ATP III	No. There was no statistically significant difference in level of physical activity between women with and without metabolic syndrome.
USA Kang, <i>et al.</i> (36)	52 women/ 39 men	Paffenbarger IPAQ	ATP III	No. Physical activity was similar between the with and without metabolic syndrome groups
Colombia Present study	89 women	IPAQ, short form	IDF	No. Lack of association between lowest physical activity level and metabolic syndrome

IPAQ: International Physical Activity Questionnaire; ATP III: National Cholesterol Education Program (NCEP) - Adult Treatment Panel  
IDF: International Diabetes Federation

Likewise, it is possible that with a larger sample size, the relationship between vigorous physical activity and clinical, biochemical and anthropometrical variables could have been evident. Furthermore, we must indicate the need for programs, including preventive and educational measures, aimed at continually promoting more active lifestyles based on the regular practice of physical activity, given that the findings of this work have revealed that moderate physical activity (estimated by minutes and days of activity) is related to the indicators of cardiovascular and metabolic risk. Likewise, development of research designs to study all the possible variables influencing the relationship between self-reported physical activity level and metabolic syndrome is also important.

### Conflicts of interest

The authors declare that there are not conflicts of interest.

### Financing

This research was supported by the Colombian Institute for Science and Technology Development (COLCIENCIAS) (Contract 1106-45921521).

### References

1. Triviño LP, Dosman V, Uribe YL, Agredo RA, Jerez AM, Ramírez-Vélez R. Estudio del estilo de vida y su relación con factores de riesgo cardiovascular en adultos de mediana edad. *Acta Med Colomb*. 2009;34:158-63.
2. Reaven GM. Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes*. 1988;37:1595-607.
3. Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med*. 2003;163:427-36. <http://dx.doi.org/10.1001/archinte.163.4.427>
4. Wildman RP, Muntner P, Reynolds K, McGinn AP, Rajpathak S, Wylie-Rosett J, *et al*. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: Prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med*. 2008;168:1617-24. <http://dx.doi.org/10.1001/archinte.168.15.1617>
5. Ford ES, Giles WH, Dietz WH. Prevalence of metabolic syndrome among US adults: Findings from the Third National Health and Nutrition Examination Survey. *JAMA*. 2002;287:356-9. <http://dx.doi.org/10.1001/jama.287.3.356>
6. Wildman RP, Muntner P, Reynolds K, McGinn AP, Rajpathak S, Wylie-Rosett J, *et al*. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: Prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med*. 2008;168:1617-24. <http://dx.doi.org/10.1001/archinte.168.15.1617>
7. Alberti KG, Zimmet P, Shaw J, IDF Epidemiology Task Force Consensus Group. The metabolic syndrome - a new worldwide definition. *Lancet*. 2005;366:1059-62. [http://dx.doi.org/10.1016/S0140-6736\(05\)67402-8](http://dx.doi.org/10.1016/S0140-6736(05)67402-8)
8. Esteghamati A, Khalilzadeh O, Rashidi A, Meysamie A, Haghazali M, Abbasi M, *et al*. Association between physical activity and metabolic syndrome in Iranian adults: National Surveillance of Risk Factors of Noncommunicable Diseases (SuRFNCD-2007). *Metabolism*. 2009;58:1347-55. <http://dx.doi.org/10.1016/j.metabol.2009.04.019>
9. Cameron AJ, Shaw JE, Zimmet PZ. The metabolic syndrome: Prevalence in worldwide populations. *Endocrinol Metab Clin North Am*. 2004;33:351-75. <http://dx.doi.org/10.1016/j.ecl.2004.03.005>
10. Hallal PC, Gomez LF, Parra DC, Lobelo F, Mosquera J, Florindo AA, *et al*. Lessons learned after 10 years of IPAQ use in Brazil and Colombia. *J Phys Act Health*. 2010;7(Suppl.2):S259-64.
11. Parra DC, McKenzie TL, Ribeiro IC, Ferreira Hino AA, Dreisinger M, Coniglio K, *et al*. Assessing physical activity in public parks in Brazil using systematic observation. *Am J Public Health*. 2010;100:1420-6. <http://dx.doi.org/10.2105/AJPH.2009.181230>
12. IPAQ group. The International Physical Activity Questionnaire, 2005. Fecha de consulta: 1 de junio de 2010. Disponible en: <https://sites.google.com/site/theipaq/>.
13. Alberti KG, Zimmet P, Shaw J. The metabolic syndrome—a new worldwide definition. *Lancet*. 2005;366:1059-62. [http://dx.doi.org/10.1016/S0140-6736\(05\)67402-8](http://dx.doi.org/10.1016/S0140-6736(05)67402-8)
14. Sarmiento OL, Schmid TL, Parra DC, Díaz-del-Castillo A, Gómez LF, Pratt M, *et al*. Quality of life, physical activity, and built environment characteristics among Colombian adults. *J Phys Act Health*. 2010;7(Suppl.2):S181-95.
15. Lobelo F, Pate R, Parra D, Duperly J, Pratt M. Burden of mortality associated to physical inactivity in Bogotá, Colombia. *Rev Salud Pública*. (Bogotá). 2006;8(Supl.2):28-41. <http://dx.doi.org/10.1590/S0124-00642006000500003>
16. Heesch KC, van Uffelen JG, Hill RL, Brown WJ. What do IPAQ questions mean to older adults? Lessons from cognitive interviews. *Int J Behav Nutr Phys Act*. 2010;7:35. <http://dx.doi.org/10.1186/1479-5868-7-35>
17. Hallal PC, Simoes E, Reichert FF, Azevedo MR, Ramos LR, Pratt M, *et al*. Validity and reliability of the telephone-administered International Physical Activity Questionnaire in Brazil. *J Phys Act Health*. 2010;7:402-9.
18. Ishii K, Shibata A, Oka K. Environmental, psychological, and social influences on physical activity among Japanese adults: Structural equation modeling analysis. *Int J Behav Nutr Phys Act*. 2010;7:61. <http://dx.doi.org/10.1186/1479-5868-7-61>
19. Blair SN, Kohl III HW, Paffenbarger RS, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: A prospective study of healthy men and women. *JAMA*. 1989;261:2395-401. <http://dx.doi.org/10.1001/jama.1989.03430170057028>
20. Farrell SW, Braun L, Barlow CE, Cheng YJ, Blair SN. The relation of body mass index, cardiorespiratory fitness, and all-cause mortality in women. *Obes Res*. 2002;10:417-23. <http://dx.doi.org/10.1038/oby.2002.58>

21. **Lee IM, Hsieh CC, Paffenbarger RS Jr.** Exercise intensity and longevity in men. The Harvard Alumni Health Study. *JAMA*. 1995;273:1179-84. <http://dx.doi.org/10.1001/jama.1995.03520390039030>
22. **Heath GW, Hagberg JM, Ehsani AA, Holloszy JO.** A physiological comparison of young and older endurance athletes. *J Appl Physiol*. 1981;51:634-40.
23. **Rodrigues ESR, Cheik NC, Mayer AF.** Nível de atividade física e tabagismo em universitários. *Rev Saúde Pública*. 2008;42:672-8. <http://dx.doi.org/10.1590/S0034-89102008000400013>
24. **Martins MD, Ricarte IF, Rocha CH, Maia RB, Silva VB, Veras AB, et al.** Blood pressure, excess weight and level of physical activity in students of a public university. *Arq Bras Cardiol*. 2010;95:192-9. <http://dx.doi.org/10.1590/S0066-782X2010005000069>
25. **Uribe YL, Triviño LP, Dosman V, Agredo RA, Jerez AM, Ramírez-Vélez R.** Asociación entre la aptitud física y los factores de riesgo de síndrome metabólico en trabajadores de una institución universitaria. *Rev Colomb Cardiol*. 2009;16:153-8.
26. **Triviño LP, Dosman V, Uribe YL, Agredo RA, Jerez AM, Ramírez-Vélez R.** Relación entre la capacidad funcional por  $VO_{2max}$  y la calidad de vida en trabajadores de una institución universitaria. *Rev Andal Med Deporte*. 2010;3:57-61.
27. **Lobelo F, Ruiz JR.** Cardiorespiratory fitness as criterion validity for health-based metabolic syndrome definition in adolescents. *J Am Coll Cardiol*. 2007;50:471-2. <http://dx.doi.org/10.1016/j.jacc.2007.04.045>
28. **Loustalot F, Carlson SA, Fulton JE, Kruger J, Galuska DA, Lobelo F.** Prevalence of self-reported aerobic physical activity among U.S. States and territories--Behavioral Risk Factor Surveillance System, 2007. *J Phys Act Health*. 2009;6(Suppl.1):S9-17.
29. **Marcondelli P, Costa THM, Schmitz BAS.** Nível de atividade física e hábitos alimentares de universitários do 3º ao 5º semestres da área da saúde. *Rev Nutr*. 2008;21:39-47. <http://dx.doi.org/10.1590/S1415-52732008000100005>
30. **American College of Sports Medicine.** Position stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in health adults. *Med Sci Sports Exerc*. 1998;30:1-34.
31. **Yu Z, Ye X, Wang J, Qi Q, Franco OH, Rennie KL, et al.** Associations of physical activity with inflammatory factors, adipocytokines, and metabolic syndrome in middle-aged and older Chinese people. *Circulation*. 2009;119:2969-77. <http://dx.doi.org/10.1161/CIRCULATIONAHA.108.833574>
32. **Esteghamati A, Khalilzadeh O, Rashidi A, Meysamie A, Haghazali M, Asgari F, et al.** Association between physical activity and insulin resistance in Iranian adults: National Surveillance of Risk Factors of Non-Communicable Diseases (SuRFNCD-2007). *Prev Med*. 2009;49:402-6. <http://dx.doi.org/10.1016/j.ypmed.2009.09.005>
33. **Abla-Mehio Sibai AM, Obeid O, Batal M, Adra N, El Khoury D, Hwalla N.** Prevalence and correlates of metabolic syndrome in an adult Lebanese population. *CVD Preven Control*. 2008;3:83-90. <http://dx.doi.org/10.1016/j.precon.2007.06.002>
34. **Dalacorte RR, Reichert CL, Vieira JL.** Metabolic syndrome and physical activity in southern Brazilian community-dwelling elders: a population-based, cross-sectional study. *BMC Public Health*. 2009;9:25. <http://dx.doi.org/10.1186/1471-2458-9-25>
35. **Delavar MA, Lye MS, Khor GL, Hassan ST, Hanachi P.** Physical activity and the metabolic syndrome in middle aged women, Babol, Mazandaran Province, Iran. *European Journal of Scientific Research*. 2008;22:411-21.
36. **Kang H, Greenson JK, Omo JT, Chao C, Peterman D, Anderson L, et al.** Metabolic syndrome is associated with greater histologic severity, higher carbohydrate, and lower fat diet in patients with NAFLD. *Am J Gastroenterol*. 2006;101:2247-53. <http://dx.doi.org/10.1111/j.1572-0241.2006.00719.x>
37. **Wannamethee G, Shaper AG.** Physical activity and stroke in British middle aged men. *BMJ*. 1992;304:597-601.
38. **Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, et al.** Walking compared with vigorous physical activity and risk of type 2 diabetes in women: A prospective study. *JAMA*. 1999;282:1433-9. <http://dx.doi.org/>